Artwork No.		Colours Us
Customer	Somex Pharma	Black
Description	Methocarbamol Tablets_750 mg_Package Leaflet	
Market	UK	
Language	English	
Size	503 x 55 mm	
Min. Font Size		
Version No.	07	
Pharma code		
Date	02.04.2022	
Software	Adobe Illustrator CS6	


INNER PAGE FRONT

STOP taking this medicine and tell your - infections, doctor immediately if you experience - mouth ulcers. any of the following rare symptoms: - mouth ulcers. - confusion, convulsions, fits, and rash or swelling of the face or neck and shortness of breath which may be a sign of an allergy. If any of the following side effects are troublesome, you should tell your doctor: If you notice any of these symptoms, you should contact your doctor: - drowsiness, difficulty in sleeping, - restlessness, anxiety or worry, - eating disorders, - wemory loss, - blurred or double vision, rapid or unus eye movements, - reddening of the eyes together with na stuffiness,

503 mm

		INNER PAGE BACK		
 WHAT METHOCARBAMOL IS AND WHAT IT IS USED FOR have ever suffered any brain damage or coma suffer from epilepsy or convulsions suffer from muscle weakness (a disease called myasthenia gravis) WAT YOU NEED TO KNOW BEFORE WAT YOU NEED TO KNOW BEFORE Warnings and precautions Talk to yous 5 doctor before taking methocarbamol if you: are allergic (hypersensitive) to methocarbamol or have had 90% (hoplems in the past with methocarbamol are allergic to any of the other ingredients of 	If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking methocarbamol. Chitdren and adolescents This medicine is not recommended for use in children due to a lack of data. 49MM	 sickness (anti-cholinergics). Psychotropic drugs for the treatment of anxiety, depression or other mental illnesses. If you are due to have anaesthetic for any reason, tell your doctor or dentist that you are taking methocarbamol. If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking methocarbamol. If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking methocarbamol. Methocarbamol Withitdod, drink and alcohol 	machinery and carry out other hazardous activities. These activities should therefore be avoided altogether until you are sure you are ont affected	them to ease the pain. Return any unused tablets to your pharmacist. If you take more Methocarbamol than you should It is important to stick to your prescribed dose. If you or someone else takes too much medicine, contact your doctor or nearest hospital emergency department immediately. Always take any medicine left over with you and also the bottle, as this will allow easier identification optime wedicine. Henext dose when it is due. Do not take a double dose of tablets to make up for a forgotten dose. If you stop taking Methocarbamol Keep taking methocarbamol until your doctor tells you to stop. If you have any tother duestions on the use of this product, ask your doctor or pharmacist. 4. POSSIBLE SIDE EFFECTS Like all medicines, methocarbamol can cause

sed	



.