

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Dapsone 50mg Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 50mg of dapsone.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Tablet.

White to off white coloured, circular shaped, biconvex uncoated tablet (5.30mm – 5.70mm) impressed with "C" on one face and "DP" on either side of a central division line on the other face.

The break line is intended for the subdivision of the tablet to provide two equal doses.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Dapsone is indicated for the treatment of the following infections in adults, adolescents and children aged 6 years or older (see section 5.1).

As part of a multi-drug regimen in the treatment of all forms of leprosy.

Treatment of dermatitis herpetiformis and other dermatoses.

Prophylaxis of malaria in combination with pyrimethamine.

Prophylaxis of *Pneumocystis carinii* pneumonia in immunodeficient subjects, especially AIDS patients.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

Adults and children over 12 years:

Multibacillary leprosy (3-drug regimen): 100mg daily for at least two years.

Paucibacillary leprosy (2-drug regimen): 100mg daily for at least six months.

Malaria prophylaxis: 100mg weekly with 12.5mg pyrimethamine.

Dermatitis herpetiformis: Initially 50mg daily, gradually increased to 300mg daily if required. Once lesions have begun to subside, the dose should be reduced to a minimum as soon as possible, usually 25-50mg daily, which may be continued for a number of years. Maintenance dosage can often be reduced in patients receiving a gluten-free diet.

Pneumocystis carinii pneumonia: In combination with trimethoprim, 50-100mg daily; 100mg twice weekly or 200mg once weekly.

Children 6-12 years:

Multibacillary leprosy (3-drug regimen): 50mg daily for at least two years.

Paucibacillary leprosy (2-drug regimen): 50mg daily for at least six months.

Elderly: Dosage should be reduced in the elderly where there is an impairment of hepatic function.

Method of Administration

For oral administration.

4.3 Contraindications

Known hypersensitivity to sulfonamides, sulfones, or any of the excipients; severe anaemia; porphyria; severe glucose-6-phosphate dehydrogenase deficiency.

4.4 Special warnings and precautions for use

Dapsone should be used with caution in patients with cardiac or pulmonary disease.

It is recommended that regular blood counts be performed during treatment with dapsone. Patients deficient in glucose-6-phosphate dehydrogenase, or methaemoglobin reductase, or with haemoglobin M are more susceptible to the haemolytic effects of dapsone.

Dapsone should be used with caution in anaemia. Severe anaemia should be treated before starting Dapsone.

Dapsone contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactose deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Excretion of dapsone is reduced and plasma concentrations are increased by concurrent administration of probenecid. Rifampicin has been reported to increase the plasma clearance of dapsone.

Increased dapsone and trimethoprim concentrations have been reported following concurrent administration in AIDS patients.

4.6 Fertility, pregnancy and lactation

Pregnancy

It is now generally considered that the benefits of dapsone in the treatment of leprosy outweigh any potential risk to the pregnant patient. Some leprologists recommend 5mg folic acid daily for leprosy patients receiving dapsone during pregnancy.

Breast-feeding

Dapsone diffuses into breast milk and there has been a report of haemolytic anaemia in a breast fed infant. While some feel that dapsone should not be used in lactating mothers, in general treatment for leprosy is continued in such patients.

Fertility

It is now generally considered that the benefits of dapsone in the treatment of leprosy outweigh any potential risk to the pregnant patient.

4.7 Effects on ability to drive and use machines

Not known.

4.8 Undesirable effects

Dapsone should be discontinued or reduced in dosage if severe lepra reactions affecting the eyes or nerve trunks occur.

Varying degrees of dose-related haemolysis and methaemoglobinaemia are the most frequently reported adverse effects of dapsone and occur in most subjects given more than 200mg daily; doses of up to 100mg daily do not cause significant haemolysis but subjects deficient in glucose-6-phosphate dehydrogenase are affected by doses above about 50mg daily. Hypoalbuminaemia and haemolytic anaemia has also been reported.

Although agranulocytosis has been reported rarely with dapsone when used alone, reports have been more common when dapsone has been used with other agents in the prophylaxis of malaria.

Rash, photosensitivity and pruritis may develop. Serious cutaneous hypersensitivity reactions occur rarely and include maculopapular rash, exfoliative dermatitis, toxic epidermal necrolysis, and Stevens-Johnson syndrome. Fixed drug eruptions have occurred.

A "dapsone syndrome" may occur after 3-6 weeks therapy; symptoms include rash, which is always present, fever, and eosinophilia. If dapsone is not stopped immediately, the syndrome may progress to exfoliative dermatitis, hepatitis, albuminuria and psychosis. Deaths have been recorded. Most patients require steroid therapy for several weeks, possibly due to the prolonged elimination time of the drug.

Peripheral neuropathy with motor loss has been reported in patients on dapsone for dermatological conditions. Peripheral neuropathy may occur as part of leprosy

reaction states and it is not an indication to discontinue dapsone. Other adverse effects occur infrequently and include anorexia, headache, hepatitis, jaundice, changes in liver function tests, insomnia, nausea, psychosis, tachycardia and vomiting.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

4.9 Overdose

Symptoms are hypoxia, methaemoglobinaemia and haemolytic anaemia.

In severe overdosage the stomach should be emptied by gastric lavage. Administration of activated charcoal by mouth has been shown to enhance the elimination of dapsone and its monoacetyl metabolite. Methaemoglobinaemia has been treated with slow IV injections of methylene blue 1-2mg/kg bodyweight, repeated after one hour if necessary. Methylene blue should not be administered to patients with glucose-6-phosphate dehydrogenase deficiency since it will not be effective. Haemolysis has been treated by infusion of concentrated human red blood cells to replace the damaged cells.

Supportive therapy includes oxygen to alleviate hypoxia, and administration of fluids to maintain renal flow and promote the elimination of dapsone.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Dapsone is a sulfone active against a wide range of bacteria.

Dapsone's mechanism of action is probably similar to that of the sulfonamides which involves inhibition of folic acid synthesis in susceptible organisms. It is usually considered to be bacteriostatic against *M leprae* although it may also possess weak bactericidal activity. It is also active against *Plasmodium* and *Pneumocystis carinii*. As with sulfonamides, antibacterial activity is inhibited by p-aminobenzoic acid.

5.2 Pharmacokinetic properties

Dapsone is almost completely absorbed from the GI tract with peak plasma concentrations occurring about 2-8 hours after a dose. Steady-state concentrations are not obtained until after at least 8 days of daily administration; doses of 100mg daily provide trough concentrations of 0.5 micrograms/ml. About 50-80% of dapsone in the circulation is bound to plasma proteins and nearly 100% of its monoacetylated metabolite is bound. Dapsone undergoes enterohepatic recycling. It is widely distributed; is present in saliva, breast milk and crosses the placenta. The half-life ranges from 10-80 hours. Dapsone is acetylated to monoacetyldapsone, the major metabolite, and other mono and diacetyl derivatives. Acetylation exhibits genetic polymorphism. Hydroxylation is the other major metabolite pathway resulting in hydroxylamine dapsone which may be responsible for dapsone-associated methaemoglobinaemia and haemolysis. Dapsone is mainly excreted in the urine, only 20% of a dose as unchanged drug.

5.3 Preclinical safety data

There are no pre-clinical data of relevance to the prescriber which are additional to that already included in other sections of the SPC.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Microcrystalline cellulose

Sodium lauryl sulphate

Croscarmellose sodium

Povidone K-30

Magnesium stearate

Purified water

6.2 Incompatibilities

None stated.

6.3 Shelf life

2 years

6.4 Special precautions for storage

Store below 25°C.

6.5 Nature and contents of container

Pack size 28's,30's

7 tablets are packed in a blister made of plain aluminium foil with white opaque PVC/PVDC foil, width 200 mm thickness.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Strandhaven Limited t/a Somex Pharma

600 High Road

Ilford, Essex,

IG3 8BS.UK

8 MARKETING AUTHORISATION NUMBER(S)

PL 15764/0136

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

26/08/2021

10 DATE OF REVISION OF THE TEXT

26/08/2021